arXiv:1207.6689v1 [hep-ph] 28 Jul 2012

Top Quark Asymmetries and Unparticle Physics

Sara Khatibi and Mojtaba Mohammadi Najafabadi

School of Particles and Accelerators,
Institute for Research in Fundamental Sciences (IPM)
P.O. Box 19395-5531, Tehran, Iran

Abstract

The measured forward-backward asymmetry (A_{FB}) in top quark pair production by the CDF and D0 collaborations has large inconsistency with the Standard Model prediction. We probe the parameters space of vector unparticle for finding an allowed region, which can explain the observed forward-backward asymmetry. In particular, we examine the necessary observables measured at the LHC and Tevatron to identify the regions in parameter space that can give top A_{FB}, the total $t\bar{t}$ cross sections and the charge asymmetry according to the LHC and Tevatron measurements. It is shown that the charge asymmetry is a powerful tool to probe the unparticle parameters space. We show that in spite of several models that proposed to explain the top A_{FB}, there exists a region in the unarticle parameters space where the top A_{FB} and the LHC charge asymmetry are satisfied simultenously.

PACS number(s): 14.65.Ha, 12.60.-i

1 Introduction

Top quark, with its mass near to the scale of electroweak symmetry breaking can be more sensitive to new physics at TeV scale than the other Standard Model (SM) partciels. Most of its properties have been examined at the Tevatron and LHC and found to be in agreement with the SM predictions [1],[2], except the observed forward-backward asymmetry in top quark pair production (A_{FB}) which has more than 2σ deviation from the SM expectation. The forward-backward asymmetry (A_{FB}) is defined as the difference between the number of top quark in the froward ($\cos\theta > 0$) and backward ($\cos\theta < 0$) region of the detector:

$$A_{FB} = \frac{N_t(\cos\theta > 0) - N_t(\cos\theta < 0)}{N_t(\cos\theta > 0) + N_t(\cos\theta < 0)} \tag{1}$$

Where θ is the top quark production angle in the $t\bar{t}$ rest frame. Recent results reported by CDF and D0 are $A_{FB} = 0.158 \pm 0.075$ [3],[4], $A_{FB} = 0.196 \pm 0.065$ [5]. While SM just predicts 0.089 [6], [7],[8],[9] for A_{FB} at loop level. There are several models that have been suggested for explaining this observation [10],[11],[12],[13],[14],[15],[16],[17],[18],[19], [20].

Since the angle θ is related to the difference of rapidities of t and $\bar{t}$ in the $p\bar{p}$ frame via:

$$y_t - y_{\bar{t}} = 2\mathrm{arctanh}(\sqrt{1 - \frac{4m_t^2}{\hat{s}}}\cos\theta) \tag{2}$$

where $y_t(y_{\bar{t}})$ is the rapidity of the top (anti)quark in the laboratory frame. The sign of $y_t - y_{\bar{t}}$ is similar to $\cos\theta$, therefore A_{FB} also can be written as:

$$A_{FB} = \frac{N_t(y_t - y_{\bar{t}} > 0) - N_t(y_t - y_{\bar{t}} < 0)}{N_t(y_t - y_{\bar{t}} > 0) + N_t(y_t - y_{\bar{t}} < 0)} \tag{3}$$

Unlike the top A_{FB}, the total $t\bar{t}$ cross section which has been measured in Tevatron is in agreement with the SM [21]:

$$\sigma_{\text{Tevatron}} = 7.56 \pm 0.63 \text{ pb.} \tag{4}$$

Note that the A_{FB} vanishes at the LHC because of the symmetric initial state. However, another observable that LHC can measure is charge asymmetry A_C. It is defined as the relative difference between top pair events with $|y_t| > |y_{\bar{t}}|$ and the events with $|y_t| < |y_{\bar{t}}|$:

$$A_C = \frac{N_t(|y_t| > |y_{\bar{t}}|) - N_t(|y_t| < |y_{\bar{t}})}{N_t(|y_t| > |y_{\bar{t}}|) + N_t(|y_t| < |y_{\bar{t}})} \tag{5}$$

At the LHC the top quarks produced in the quark-antiquark annihilation process are statistically more boosted to the beam direction in comparison with the antitop quark. This is because of the fact that the top quark prefers to fly in the direction of the incident quark which carries a larger longitudinal momentum. As a consequence, a charge asymmetry as described above is generated.

The ATLAS and CMS measurements for the charge asymmetry are: $A_C = -0.018 \pm 0.036$ [22], $A_C = -0.013 \pm 0.041$ [23], and the SM prediction is $A_C = 0.0115$ [9]. It is clear that within the uncertainties the standard model prediction is in agreement with the measured values at the LHC.

It is notable that some of the SM extensions proposed to explain the Tevtatron A_{FB} also predict sizeable charge asymmetry at the LHC [24],[25]. Therefore, in those models there exists a tension between the top forward-backward asymmetry at Tevatron and the LHC charge asymmetry. From another side, the LHC charge asymmetry measurement is consistent with the SM expectation consequently the models which predict also enhancement in A_C are disfavored. For example, it has been shown that in the W' and Z' models there is a tight correlation between A_{FB} and A_C so that the measured charge asymmetry excludes these models [24],[26],[27],[28]. In [29], the effective Lagrangian approach has been utilized to explain the A_{FB}. In this approach an enhancement in A_C is also expected, in particular at large $t\bar{t}$ invariant mass region. It has been shown in [26] that there is an apparent tension between the forward-backward asymmetry and the charge asymmetry in axigluon model.

It seems difficult to develop a model that can produce large A_{FB} deviated from the SM prediction according to Tevatron measurement but A_C is consistent with the SM value. There are studies on this issue which for example can be found in [24],[25].

The measured top pair cross section at the LHC confirms the SM expectation [30]:

$$\sigma_{\rm LHC} = 165 \pm 13.3 \text{ pb}. \tag{6}$$

As we know the scale invariance is broken in the SM by the mass of particles. However, one can imagine the existence of a scale invariant sector at TeV scale or higher that might be observed at the LHC or future colliders. In [31], [32], Georgi has suggested a model known as unparticle in which there is a nontrivial sector that is exactly scale invariant. The SM fields have very weak interactions with the scale invariant sector via non-renormalizable effective interactions.

Since particles have definite nonzero masses, scale invariant stuff is made of unparticles. Note that no signal of this sector has been observed yet and if it exists, evidence of unparticle could show up e.g. as large missing energy, change in the production cross sections branching ratios, and different behavior in angular distributions of the final state particles at colliders [33].
Top physics could be affected by the new interactions with unparticles. There have been already several studies to explain the top A_{FB} through unparticle. In [34], vector unparticles that carry color charges are used to explain the observed forward-backward asymmetry. Recently, in [35], the contribution of unparticle to A_{FB} and spin correlation in $t\bar{t}$ production at the Tevatron has been investigated.

In this work, we study the effects of color singlet vector unparticles on the forward-backward asymmetry and charge asymmetry at Tevatron and LHC, respectively. In particular, we investigate the tension between A_C and A_{FB} and perform a full scan on the main unparticle parameters space.

The organization of this letter is as follows. Next section is devoted to unparticle physics and its effect on top prodution rate. In section 3, we show our numerical calculations and discuss the results. Finally, conclusions are presented in section 4.

2 Influence of unparticle on top pair production

The effects of unparticle on top properties at hadron colliders have been intensively studied in the literatures [36],[37], [38], [39], [40], [41], [42], [43]. Also, there are some papers in which the top A_{FB} at the Tevatron has been studied. In [34], the authors have found the regions of parameters where colored vector unparticle can produce the values of top A_{FB} and the top pair cross sections compatible with the Tevatron measurements. In [35], the influence of vector and tensor unparticle, including color, on top pair cross section and the forward-backward asymmetry has been investigated.
Effective interaction of vector unparticle with SM fields are given as follows [33]:

$$\lambda_1 \frac{1}{\Lambda^{d_{\mathcal{U}}-1}} c_v \bar{f} \gamma_\mu f O^\mu_{\mathcal{U}} \quad , \quad \lambda_1 \frac{1}{\Lambda^{d_{\mathcal{U}}-1}} c_a \bar{f} \gamma_\mu \gamma_5 f O^\mu_{\mathcal{U}} \tag{7}$$

Where λ_1 is dimensionless effective couplings labeling vector unparticle operator. The coefficients c_v, c_a represent vector and axial vector couplings of vector unparticle, respectively. The parameter $d_{\mathcal{U}}$ is the scaling dimension of the unparticle operators and Λ denotes the effective mass scale above which unparticle is formed.

Within the SM at hadron colliders, $t\bar{t}$ pairs are produced either via quark-antiquark annihilation or through gluon-gluon fusion. With considering new interactions of vector unparticle with SM fields, only the partonic cross section for $t\bar{t}$ production via quark-antiquark annihilation is modified, because vector unparticle only interacts with fermionic fields and it dosen't couple to gluons.

The parton level differential cross section for the process of $q\bar{q} \to t\bar{t}$ at leading order in the presence of color singlet vector unparticle is as follows [36]:

$$\begin{aligned} \frac{d\hat{\sigma}}{d\hat{t}}(q\bar{q} \to t\bar{t}) &= \frac{A_V^2}{8\pi\hat{s}^2(\hat{s})^{4-2d_{\mathcal{U}}}}\Big[c_a^4(2m^4 - 4(\hat{s}+\hat{t})m^2 + (\hat{s}+\hat{t})^2 + \hat{t}^2) \\ &+ c_v^4(2m^4 - 4\hat{t}m^2 + (\hat{s}+\hat{t})^2 + \hat{t}^2) \\ &+ 2c_v^2c_a^2(2m^4 - 2(3\hat{s}+2\hat{t})m^2 + 3\hat{s}^2 + 2\hat{t}^2 + 6\hat{s}\hat{t})\Big] \\ &+ \frac{d\sigma^0_{q\bar{q}}}{d\hat{t}} \end{aligned} \tag{8}$$

where

$$A_V = \frac{\lambda_1^2 A_{d_{\mathcal{U}}}}{2\sin(d_{\mathcal{U}}\pi)\Lambda^{2(d_{\mathcal{U}}-1)}}, \tag{9}$$

and

$$A_{d_{\mathcal{U}}} = \frac{16\pi^2\sqrt{\pi}}{(2\pi)^{2d_U}}\frac{\Gamma(d_{\mathcal{U}}+\frac{1}{2})}{\Gamma(d_{\mathcal{U}}-1)\Gamma(2d_{\mathcal{U}})}. \tag{10}$$

In the cross sections relation, $\frac{d\sigma^0_{q\bar{q}}}{d\hat{t}}$ is the SM part which has the following form:

$$\frac{d\sigma^0_{q\bar{q}}}{d\hat{t}} = \frac{4\pi\alpha_s^2}{9\hat{s}^4}\Big[2m^4 - 4\hat{t}m^2 + (\hat{s}+\hat{t})^2 + \hat{t}^2\Big]. \tag{11}$$

In Eq.8, for the case that $c_v = 1$ is corresponding to vector unpartcile, $c_v = c_a = 1$ is corresponding to vector unparticle with right-handed coupling to the SM fields and $c_v = -c_a = 1$ presents the vector unparticle with left-handed coupling. According to Eq. 8, the cross section is similar in both cases with $c_v = c_a = 1$ and $c_v = -c_a = 1$. Therefore, the $t\bar{t}$ cross section and forward-backward asymmetry in this scenario are chirality independent or blind to left-hand or right-handed couplings.

3 Numerical Results and Discussion

The hadronic cross section is obtained from the parton level cross section convoluted with the parton distribution functions according to the following relation:

$$d\sigma(pp(\bar{p}) \to t\bar{t}) = \sum_{ij} \int_0^1 dx_1 \int_0^1 dx_2 f_i(x_1, Q^2) f_j(x_2, Q^2) d\hat{\sigma}_{ij}. \tag{12}$$

where $f_{i,j}$ are the parton distribution functions. In the numerical calculations, the top quark mass has been set $m_t = 173$ GeV. All cross sections at the partonic level is calculated by employing CTEQ6 parton distribution functions [44] with a choice of $Q^2 = m_t^2$. The calculation is performed at fixed renormalization and factorization scale $\mu_R = \mu_F = m_t$. We present our numerical result at Tevatron with $\sqrt{s} = 1.96$ TeV and at LHC with $\sqrt{s} = 7$ TeV. The strong couling constant α_S at the scale of top mass is set to 0.11. Indeed, the cross sections that we get from the calculations are the leading order values. Therefore, we scale the tree-level calculation by a k-factor of 1.3, so that the leading order calculations match with the higher order results for the case of $m_t = 173$ GeV/c^2. The NNLO cross section of top pair production at Tevatron is 7.08 pb and 163 pb at the LHC with the center-of-mass energy of 7 TeV [45].

As we mentioned before, the results are chirality independent and the right-handed and left-handed unparticle couplings to the SM fields give similar cross sections and asymmetries in top pair events. In the case of having pure vector unparticle i.e. $c_v = 1$ and $c_a = 0$, negligible forward-backward asymmetry is produced which can not compensate the observed value by Tevatron experiments.

First, we present all observables in terms of $d_{\mathcal{U}}$ for three various values of Λ, and consider $\lambda_1 = 1$ and $c_a = c_v = 1$. Then we identify an allowed region in the $d_{\mathcal{U}}$, Λ plane for all observables in the same plot. We concentrate on the values of unparticle parameters which are physically interesting, i.e. $1 < d_{\mathcal{U}} < 2$ and Λ at the order of few TeV.

The forward-backward asymmetry (A_{FB}) at Tevatron and the charge asymmetry at the LHC are shown in Fig. 1. The shaded area is the experimental band. As it can be seen, for a specific value of $d_{\mathcal{U}}$ the forward-backward asymmetry grows when Λ decreases, i.e. unparticle can produce larger asymmetry by assuming small values of Λ. Note that for larger values of Λ, the allowed interval of $d_{\mathcal{U}}$ parameter that can produce desirable forward-backward asymmetry becomes smaller. According to Fig. 1 at $\Lambda = 1$ TeV, unparticle with any value of $d_{\mathcal{U}}$ in the



Figure 1: *Left: The forward-backward asymmetry* (A_{FB}) *in top pair production generated by vector unparticle for* $\lambda_1 = 1$ *and* $c_a = c_v = 1$ *and various values of* Λ. *Shaded region is the band measured by the CDF experiment. Right: The charge asymmetry at the LHC with the same parameters as* A_{FB} *plot and different values of* Λ.

range of 1.2 to 1.32 can generate the desired A_{FB}.

The behavior of charge asymmetry is different from A_{FB}. The charge asymmetry A_C grows with increasing $d_{\mathcal{U}}$, reaches to a maximum value at $d_{\mathcal{U}} = 1.1$ then it decreases and tends to the SM expectation at $d_{\mathcal{U}}$ in the tail. Since the peak position does not move for various values of Λ. According to Fig.1, charge asymmetry is a powerful observable in scanning the parameter space of unparticle. The shaded region is according to the CMS measurement. For example, when $\Lambda = 1$ TeV, unparticle with $d_{\mathcal{U}} = 1.28$ is excluded. For larger values of Λ, the exclusions interval is smaller. The combination of limits from A_C and the allowed band for A_{FB} leads to the allowed interval of 1.28 to 1.32 for $d_{\mathcal{U}}$ at the value of $\Lambda = 1$ TeV. This interval gets smaller when we move to higher values of Λ.

The total cross section of top pair production at the Tevatron and LHC versus $d_{\mathcal{U}}$ for various values of Λ are shown in Fig. 2. The bands are according to the present Tevatron and LHC measurements. There is a small interval where the cross sections are sensitive to $d_{\mathcal{U}}$. This interval is reduced when we go to high values of Λ. For example, for both LHC and Tevatron, when $\Lambda = 1$ TeV, the cross sections are tending downwards and become compatible with the SM predictions at $d_{\mathcal{U}} = 1.4$. For $\Lambda = 5$ TeV, the position where the cross sections become equal to their SM values is around $d_{\mathcal{U}} = 1.2$.

In Fig.3, we present the allowed regions in the plane of $(d_{\mathcal{U}}, \Lambda)$ which satisfy the cross section



Figure 2: *The total cross cection of top pair production by vector unparticle for* $\lambda_1 = 1$ *and* $c_a = c_v = 1$ *at the Tevatron (left) and LHC (right). Shaded regions are the experimental bands.*

of $t\bar{t}$ at Tevatron and LHC, the measured forward-backward asymmetry by Tevatron and the charge asymmetry. The area between two dashed black curves consists of the points that can produce the observed forward-bakward asymmetry by Tevatron. The right side of the green dashed-dotted curve is the allowed region according to the charge asymmetry measurement of LHC. The right sides of blue and red curves are satisfied by the Tevatron and LHC cross section measurements.

As it can be seen in Fig.3, the band in which A_{FB} can be generated consistent with the Tevatron observation gets narrower with increasing the value of Λ. The limit that charge asymmetry applies on the parameter space is almost the same as the limit coming from the LHC cross section. The allowed region arisng from the top pair cross section at the Tevatron is tigher than the LHC. Finally, the region compatible with all measurements considered in this study has been shown in Fig.3 by shaded area between the solid blue curve and the dashed black curve in the right side. When we move toward the large values of Λ, the allowed area in the parameter space which can produce the desired forward-backward asymmetry gets smaller.

As mentioned before the relation between the Tevatron A_{FB} and the LHC charge asymmetry A_C is model dependent. For example, some models [26],[27] can generate the forward-backward asymmetry but the LHC charge asymmetry excludes the the regions where A_{FB} is genretaed according to the Tevatron measurememts. Contrary to models like the W',Z', axigluon models in which there is a tight tension between A_{FB} and A_C, color singlet vector unparticle (with left or right handed couplings) can generate large forward-backward asymmetry keeping the cross



Figure 3: *Region of* Λ *(in GeV) in terms of* d_u *consistent with Tevatron measurements of the* $t\bar{t}$ *forward-backward asymmetry (region between two dashed black curves). The right side of the solid red (blue) curves is consistent with the LHC (Tevatron) cross section. The consistent region with the LHC charge asymmetry is the area in the right side of the dot-dashed green curve. Shaded region is the allowed region according to all measurements.*

sections and charge asymmetry consistent with the SM expectations.

4 Conclusions

We investigated whether there is an allowed region where the top forward-backward asymmetry measured by Tevatron experiments can be produced by color singlet vector unparticles. Then, we showed that contrary to some models in which there is a tight tension between A_{FB} and A_C, color singlet vector unparticle (with left or right handed couplings) can generate large forward-backward asymmetry keeping the cross sections and charge asymmetry consistent with the measurements and of course with the SM expectations. We found an area in the plane of $(d_{\mathcal{U}}, \Lambda)$ which gives the observed forward-backward asymmetry, the $t\bar{t}$ cross sections and the charge asymmetry compatible with the experimental measurements. The allowed interval of $d_{\mathcal{U}}$ becomes smaller when we move up to the larger values of Λ. Moreover, we found that pure vector unparticle can not generate large enough forward-backward asymmetry.

References

[1] F. -P. Schilling, Int. J. Mod. Phys. A **27**, 1230016 (2012) [arXiv:1206.4484 [hep-ex]].

[2] W. Bernreuther, arXiv:0805.1333 [hep-ph].

[3] T. Aaltonen et al. [CDF Collaboration], Phys. Rev. D 83, 112003 (2011).

[4] The CDF Collaboration, CDF Note 10807.

[5] V. Abazov et al. [D0 Collaboration], Phys. Rev. Lett. 84, 112005 (2011), arXiv:1107.4995 [hep-ex].

[6] P. Ferrario and G. Rodrigo, Phys. Rev. D 78, 094018 (2008).

[7] W. Bernreuther and Z. G. Si, Nucl. Phys. B 837, 90 (2010).

[8] W. Hollik and D. Pagani, arXiv:1107.2606 [hep-ph].

[9] J. H. Kuhn and G. Rodrigo, JHEP 1201, 063 (2012).

[10] P. Ferrario and G. Rodrigo, Phys. Rev. D 80, 051701 (2009) [arXiv:0906.5541 [hep-ph]].

[11] J. A. Aguilar-Saavedra and M. Perez-Victoria, Phys. Lett. B 705, 228 (2011) [arXiv:1107.2120 [hep-ph]]; G. M. Tavares and M. Schmaltz, Phys. Rev. D 84, 054008 (2011) [arXiv:1107.0978 [hep-ph]].

[12] S. Jung, H. Murayama, A. Pierce and J. D. Wells, Phys. Rev. D 81, 015004 (2010) [arXiv:0907.4112 [hep-ph]].

[13] K. Blum, Y. Hochberg and Y. Nir, JHEP 1110, 124 (2011) [arXiv:1107.4350 [hep-ph]].

[14] J. Shu, T. M. P. Tait and K. Wang, Phys. Rev. D 81, 034012 (2010) [arXiv:0911.3237 [hep-ph]].

[15] R. Barcelo, A. Carmona, M. Chala, M. Masip and J. Santiago, arXiv:1110.5914 [hep-ph].

[16] R. Barcelo, A. Carmona, M. Masip and J. Santiago, arXiv:1106.4054 [hep-ph].

[17] R. Barcelo, A. Carmona, M. Masip and J. Santiago, Phys. Rev. D **84** (2011) 014024 [arXiv:1105.3333 [hep-ph]].

[18] B. Bhattacherjee, S. S. Biswal and D. Ghosh, Phys. Rev. D **83**, 091501 (2011) [arXiv:1102.0545 [hep-ph]].

[19] B. Grinstein, C. W. Murphy, D. Pirtskhalava and P. Uttayarat, arXiv:1203.2183 [hep-ph].

[20] K. Cheung, W. -Y. Keung and T. -C. Yuan, Phys. Lett. B **682**, 287 (2009) [arXiv:0908.2589 [hep-ph]].

[21] D0 Collaboration, Phys. Lett. B 704, (2011) 403-410.

[22] G. Aad et al. [ATLAS Collaboration], arXiv:1203.4211 [hep-ex].

[23] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **709**, 28 (2012) [arXiv:1112.5100 [hep-ex]]; The CMS Collaboration, CMS PAS TOP-11-030.

[24] J. A. Aguilar Saavedra and M. Perez-Victoria, Phys. Rev. **D 84**, (2011) 115013.

[25] J. A. Aguilar Saavedra and A. Juste, arXiv:1205.1898.

[26] S. Fajfer, J. F. Kamenik and B. Melic, arXiv:1205.0264 [hep-ph].

[27] S. Y. Ayazi, S. Khatibi and M. Mohammadi Najafabadi, arXiv:1205.3311 [hep-ph].

[28] S. Y. Ayazi, arXiv:1207.0643 [hep-ph].

[29] C. Delaunay, O. Gedalia, Y. Hochberg and Y. Soreq, arXiv:1207.740 [hep-ph].

[30] The CMS Collaboration, CMS-PAS-TOP-11-024.

[31] H. Georgi, Phys. Rev. Lett. **98**, 221601 (2007) [hep-ph/0703260].

[32] H. Georgi, Phys. Lett. B **650**, 275 (2007) [arXiv:0704.2457 [hep-ph]].

[33] K. Cheung, W. -Y. Keung and T.-C. Yuan, Phys. Rev. D **76** (2007) 055003; L. -G. Bian, arXiv:1108.1538 [hep-ph]; T. Kikuchi and N. Okada, Phys. Lett B **661**, 360 (2008); K. Cheung, W. -Y. Keung and T. -C. Yuan, Phys. Rev. Lett. **99**, 051803 (2007); S. -L. Chen and X. -G. He, Phys. Rev. D **76**, 091702 (2007) [arXiv:0705.3946 [hep-ph]].

[34] C. -H. Chen, G. Cvetic and C. S. Kim, Phys. Lett. B **694**, 393 (2011) [arXiv:1009.4165 [hep-ph]].

[35] M. Dahiya, S. Dutta and R. Islam, arXiv:1206.5447 [hep-ph].

[36] A. T. Alan and N. K. Pak, Europhys. Lett. **84**, 11001 (2008) [arXiv:0708.3802 [hep-ph]].

[37] D. Choudhury and D. K. Ghosh, Int. J. Mod. Phys. A **23**, 2579 (2008) [arXiv:0707.2074 [hep-ph]].

[38] M. Arai, N. Okada and K. Smolek, Phys. Rev. D **79**, 074019 (2009) [arXiv:0902.0418 [hep-ph]].

[39] A. T. Alan, N. K. Pak and A. Senol, Europhys. Lett. **83**, 21001 (2008) [arXiv:0710.4239 [hep-ph]].

[40] H. -F. Li, H. -l. Li, Z. -G. Si and Z. -J. Yang, Commun. Theor. Phys. **51**, 707 (2009) [arXiv:0802.0236 [hep-ph]].

[41] I. Sahin, Eur. Phys. J. C **60**, 431 (2009) [arXiv:0802.2818 [hep-ph]].

[42] T. M. Aliev, O. Cakir and K. O. Ozansoy, Phys. Lett. B **670**, 336 (2009) [arXiv:0809.2327 [hep-ph]].

[43] T. M. Aliev and K. O. Ozansoy, arXiv:0905.1597 [hep-ph].

[44] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky and W. K. Tung, JHEP **0207**, 012 (2002) [arXiv:hep-ph/0201195].

[45] N. Kidonakis, Phys. Rev. D **82**, 114030 (2010) [arXiv:1009.4935[hep-ph]].